

Stenoma cymogramma, n. sp.

♂. 18-19 mm. Head white, back of crown tinged grey. Palpi white, a dark grey streak on basal $\frac{2}{3}$ of second joint, extreme base of terminal joint dark grey. Thorax light grey, shoulders narrowly whitish. Forewings slightly narrowed posteriorly, apex rounded, termen somewhat obliquely rounded; 2 from near angle, 3-5 approximated, 7 to costa; light brownish-grey, paler or whitish-tinged towards costa anteriorly, costal edge white; three somewhat curved oblique dark fuscous lines, first two irregular, first from towards costa at $\frac{1}{4}$ to dorsum beyond middle, thickened dorsally, second from middle of costa to dorsum at $\frac{4}{5}$, second discal stigma perceptible on this, third from costa at $\frac{3}{4}$ rather abruptly curved on lower half to dorsum just before tornus; eight blackish marginal dots round posterior part of costa and termen: cilia whitish-grey, a grey subbasal shade. Hindwings 3 and 4 rather short-stalked; pale grey, whitish-tinged anteriorly; costa dilated on basal $\frac{2}{3}$, with fringe of white scales, a yellowish subcostal groove from base to middle; cilia whitish, a pale grey subbasal line. Forewings beneath with long ochreous-whitish expansible subdorsal hairpencil from base lying in subcostal groove of hindwings.

PERU, R. Napo, Iquitos, March to May (*Parish*); 2 ex.

Stenoma neocrossa, n. sp.

♂. 13 mm. Head ochreous-white. Palpi white, basal half and subapical ring of second joint, base and oblique suprmedian ring of terminal joint dark fuscous. Thorax ochreous-white, suffused light grey posteriorly. Forewings suboblong, apex rounded, termen somewhat obliquely rounded; 2-5 closely approximated at base, 6 and 7 nearly approximated throughout, 7 indefinite; white, dorsal half tinged brownish; a fine dark fuscous median dash near base; stigmata moderate, dark fuscous, plical rather obliquely beyond first discal, a short oblique streak above and before first discal pointing to it; an undefined patch of fuscous suffusion extending along anterior half of dorsum and touching plical stigma; oblique transverse dark fuscous lines from costa at middle and $\frac{3}{4}$, first straight, irregular, interrupted in disc, nearly preceding second discal stigma, second curved on lower half, dorsal half between these forming a fuscous blotch becoming dark fuscous anteriorly; three præmarginal black dots on apical part of costa, lowest largest, and two or three minute and undefined on termen: cilia white. Hindwings 3 and 4 moderately stalked; ochreous-whitish, suffused light grey posteriorly; a fringe of long raised ochreous-whitish hairs along lower margin of cell and basal half of vein 2; cilia whitish.

PERU, R. Napo, May (*Parish*); 1 ex.

Stenoma isoplintha, n. sp.

♂ ♀. 22-23 mm. Head white, crown pale grey except on sides. Palpi white, a streak on basal half of second joint, and base of terminal joint dark grey. Thorax grey, shoulders whitish. Fore-

wings oblong, apex rounded-obtuse, termen nearly vertical; 2-5 approximated at base, 7 to apex; white, in ♀ tinged ochreous except on costa; dorsal half ochreous-whitish, mainly occupied by three subquadrate dark grey blotches; in ♀ undefined longitudinal streaks of greyish suffusion anteriorly beneath costa and above middle; a better-marked longitudinal grey streak above anterior part of second dorsal blotch, shorter in ♂; a very oblique dark grey dash above end of cell, and a slight mark preceding it; a nearly straight dark grey line from beneath costa at $\frac{3}{5}$ to posterior angle of third dorsal blotch; in ♂ 3, in ♀ 6 dark grey præmarginal dots round apex and termen, largest about apex: cilia snow-white, basal third light grey on termen. Hindwings 3 and 4 short-stalked; light grey; cilia light grey, on costa white.

BRAZIL, Parintins, October (*Parish*); 2 ex.

Stenoma tetrapetra, n. sp.

♀. 25 mm. Head white. Palpi white, a grey streak on basal half of second joint. Thorax white, tinged grey posteriorly. Apex of abdomen yellowish. Forewings oblong, apex rounded-obtuse, termen rounded, little oblique; 2-5 approximated at base, 7 to apex; white; four rather dark grey blotches occupying most of dorsum, not nearly reaching half across wing, first elongate, second and third subquadrate, fourth small, triangular, tornal; an indistinct greyish streak above middle from near base to $\frac{1}{4}$, and a stronger one in disc from $\frac{1}{5}$ to $\frac{2}{5}$; second discal stigma dark grey, three greyish longitudinal marks above cell forming a very oblique series pointing to this; a short fine grey dash beyond cell above middle; five minute dark grey dots on termen: cilia white. Hindwings 3 and 4 short-stalked; ochreous-whitish suffused pale grey on posterior half; cilia white.

BRAZIL, Teffé, December (*Parish*); 1 ex.

Stenoma cremastis, n. sp.

♀. 24 mm. Head white, crown light grey except sides. Palpi white, a light greyish streak on basal half of second joint. Thorax grey, shoulders whitish. Forewings oblong, apex rounded, termen rounded, little oblique; 2-5 approximated at base, 7 to costa (indefinite); white, dorsal half mostly suffused pale ochreous-grey, somewhat darker towards dorsum anteriorly; three suffused dark grey dorsal blotches, not reaching half across wing, first subtriangular, before middle of wing, second subquadrate, about $\frac{3}{4}$, third small, triangular, tornal; a small dark grey spot in disc at $\frac{1}{4}$; second discal stigma dark fuscous, a short oblique fuscous mark above cell before this; a faint interrupted incurved fuscous line from costa at $\frac{4}{5}$ to apex of third dorsal spot; 5 or 6 dark fuscous marginal dots round apex and termen: cilia white, on tornus tinged greyish. Hindwings 3 and 4 in Jurimaguas specimen coincident (probably an abnormality), in Manaus specimen short-stalked; pale grey, rather darker posteriorly; cilia whitish-grey, round apex white.

PERU, Jurimaguas, March (type); BRAZIL, Manaus, November (*Parish*); 2 ex. These agree so well, especially in the peculiar and very characteristic form of the posterior line of forewings, that I regard them as identical, notwithstanding the different neuration of hindwings.

***Stenoma subdulcis*, n. sp.**

♂. 16 mm. Head white. Palpi white, a dark fuscous streak on basal half of second joint. Thorax very pale grey, shoulders white. Forewings oblong, costa nearly straight posteriorly, apex obtuse-rectangular, termen vertical, nearly straight; 2 from towards angle, 7 to termen; pale grey; a broad white costal band from base to near apex, posteriorly pointed, costal edge ochreous-tinged, below band a streak of whitish-ochreous suffusion to end of cell; an inwards-oblique irregular dark fuscous streak from dorsum beyond middle to fold, some slight suffusion about fold before and beyond this; a waved ochreous-whitish terminal line: cilia white. Hindwings 3 and 4 almost connate, 6 and 7 nearly approximated at base; light grey; cilia white.

BRAZIL, Santarem, August (*Parish*); 1 ex.

***Stenoma chloromis*, Meyr.**

A fresher example from La Cambre, Honduras, is 29 mm., with head light violet-grey, thorax yellow-ochreous, forewings with grey area extended to costa on basal fourth, leaving a broad white costal band beyond this.

***Stenoma remorsa*, n. sp.**

♀. 30 mm. Head white. Palpi white, a streak on basal half of second joint, and base of terminal joint dark fuscous. Thorax whitish-ochreous, a quadrate blackish dorsal spot. Forewings oblong, slightly expanded posteriorly, costa almost straight, apex rectangular, termen almost straight, vertical; 2 from towards angle, 7 to termen; whitish-ochreous; an oval blackish blotch along anterior half of dorsum; a triangular blackish blotch extending in disc from $\frac{1}{3}$ to $\frac{2}{3}$, separated from preceding by an oblique curved streak of groundcolour, and connected with dorsum by an oblique blackish streak beyond this; four blackish interneural streaks in costal area; on terminal area a band formed of confluent interneural streaks of grey suffusion, not reaching discal blotch or termen: cilia whitish-ochreous, basal half grey. Hindwings 3 and 4 connate, 6 and 7 nearly approximated towards base; light grey, rather darker towards apex; cilia whitish-ochreous, round apex suffused light grey.

BOLIVIA, R. Suruta, Dept. Santa Cruz, 1300 feet; 1 ex.

***Stenoma neoptila*, n. sp.**

♀. 25 mm. Head ochreous-white. Palpi white, second joint whitish-ochreous except tip. Thorax whitish-ochreous. Forewings somewhat dilated posteriorly, apex rounded, termen rounded, little

oblique; 2 from very near angle, 7 to apex (indefinite); white, costal edge whitish-ochreous; a dark fuscous patch extending along anterior half of dorsum but leaving dorsum slenderly ochreous-white, reaching nearly half across wing, at base touching costa, immediately beyond it a pale ochreous triangular dorsal blotch, followed by an inwards-oblique broad dark fuscous fasciate blotch from tornus reaching $\frac{3}{4}$ across wing, suffused posteriorly and above: cilia white, towards tornus whitish-ochreous suffused grey. Hindwings 3 and 4 connate, 6 and 7 approximated towards base; yellow-whitish; cilia yellow-whitish, on costa about $\frac{2}{3}$ a broad patch of long projecting grey scales.

BRAZIL, R. Trombetas, September (*Parish*); 1 ex.

***Stenoma adytodes*, n. sp.**

♂. 26-28 mm. Head white, sides of forehead fuscous. Palpi white or whitish, second joint fuscous. Thorax white, shoulders dark fuscous. Forewings elongate, rather dilated, apex obtuse, termen slightly rounded, somewhat oblique; 2-5 approximated, 7 to termen; dark fuscous; a broad white costal streak from near base to beyond middle, ends pointed, costal edge pale ochreous; an irregular-edged rounded white apical blotch extending on costa to $\frac{2}{3}$; a slender pale brownish-ochreous dorsal streak throughout, expanded into a triangular postmedian blotch tipped white, tornal extremity also white: cilia round tornus pale brownish-ochreous, round apical blotch white, between these grey. Hindwings 3 and 4 connate, 6 and 7 connate; whitish-ochreous suffused light grey, apex whitish; cilia concolorous, on costa at $\frac{2}{3}$ a thickened patch of ochreous-grey scales.

PERU, Oconeque, Carabaya, 7000 feet, February (*Ockenden*); 3 ex.

***Stenoma leucosaris*, n. sp.**

♂. 20 mm. Head whitish. Palpi white, second joint lilac-brown. Thorax light lilac-ochreous, shoulders whitish. Forewings suboblong, apex rounded, termen vertical; 2-5 approximated, 8 to apex (indefinite); light lilac-ochreous; costal edge very finely whitish-ochreous, towards base fuscous; markings rather dark purplish-fuscous; stigmata moderate, plical very obliquely beyond first discal; a small mark on costa at $\frac{1}{3}$; a dot towards costa before middle; flattened-triangular spots on costa about middle and $\frac{3}{4}$, from first an interrupted shade strongly curved round beyond second discal stigma, thence nearly straight to dorsum about $\frac{2}{3}$ but nearly obsolete on lower portion, from second a transverse series of dots excurved in disc, rather indented above and sinuate below this, running to dorsum near tornus; a terminal series of cloudy dots obscurely coppery-tinged: cilia pale glossy violet-grey. Hindwings costa somewhat expanded, 3 and 4 connate; ochreous-whitish, an undefined terminal band of light grey suffusion; cilia whitish, round apex and upper half of termen a grey subbasal shade. Forewings beneath with long whitish fringe on 1a from base reaching end of cell.

PERU, Papitinga, June (*Parish*); 1 ex. Extremely similar to *siraphora*, but apex of forewings more rounded, subdorsal fringe in *siraphora* yellowish and reaching to beyond end of cell, hindwings more suffused grey.

***Stenoma clysmographa*, n. sp.**

♂. 30 mm. Head whitish-brown. Palpi whitish, second joint suffused light brownish. Thorax light brownish. Forewings rather dilated, apex obtuse-rectangular, termen straight, vertical; 2 from towards angle, 8 to termen; without subdorsal fringe beneath; brown; markings suffused, pale greyish, viz., an irregular blotch towards costa before middle, a roundish spot about fold beneath this, two small dots longitudinally placed in disc beyond middle, some slight irroration towards costa about $\frac{2}{3}$, a curved subterminal shade from above middle to dorsum, thickened in disc and with waved posterior edge, and a terminal streak becoming obsolete towards extremities; a terminal row of dark fuscous dots: cilia pale brownish-grey. Hindwings 3 and 4 very short-stalked; dark fuscous; cilia grey, suffused darker towards base.

***Stenoma phaeomystis*, n. sp.**

♂. 27 mm. Head brownish, some whitish tinge on forehead. Palpi whitish, second joint externally and terminal joint anteriorly suffused brownish. Antennæ scaled dorsally towards middle, ciliations under 1. Thorax dark brown, patagia brown suffused brownish-ochreous anteriorly. Abdomen blackish. Forewings rather dilated posteriorly, apex obtuse, termen slightly sinuate, somewhat oblique; 2 from very near angle, 8 to apex; violet-fuscous, becoming ochreous-brown towards costa posteriorly, and dark purple-brown towards dorsum and termen. dorsal edge suffused blackish: cilia brownish, base pale ochreous-brown on upper part of termen. Hindwings 3 and 4 connate, 6 and 7 stalked; grey irrorated blackish hairscales; cilia light grey, a darker subbasal line.

PERU, R. Ucuyali; 1 ex.

***Stenoma capnocrossa*, n. sp.**

♂. 25 mm. Head, thorax pale brownish-ochreous, face whitish-ochreous. Palpi ochreous-whitish, second joint pale ochreous infuscated externally except apex. Forewings moderate, somewhat dilated, apex obtuse-rectangular, termen straight, vertical; 2 from near angle, 8 to termen; light brownish-ochreous; plical and second discal stigmata very small, dark fuscous: cilia whitish-ochreous, on termen rather dark fuscous. Hindwings 3-5 nearly approximated at base; rather dark grey; cilia whitish-grey.

BRAZIL, Tefic, January (*Parish*); 1 ex.

***Stenoma myrodora*, n. sp.**

♂ ♀. 21-22 mm. Head, thorax light greyish-ochreous, face ochreous-whitish. Palpi ochreous-grey-whitish, an obscure

purplish-grey streak on basal half of second joint. Forewings slightly dilated, apex rounded-obtuse, termen rounded, vertical; 2 from near angle, 8 to apex (very indefinite); greyish-ochreous; three obscure rather irregular grey transverse lines, first rather curved, rather oblique, lower half forming posterior limit of a triangular purple dorsal blotch, somewhat marbled with ground-colour, second and third obtusely angulated in middle, space between these forming a suffused purplish blotch on dorsal half; second discal stigma forming a dark grey transverse mark preceding second line; a fuscous terminal line: cilia pale grey. Hindwings 3 and 4 connate; grey; cilia pale grey. Thorax ♂ with a moderately long whitish hairpencil lying along side from beneath base of forewings.

BRAZIL, Teffé, January (*Parish*); FRENCH GUIANA, R. Maroni; 4 ex.

***Stenoma endochra*, n. sp.**

♂. 23 mm. Head, thorax brownish-ochreous, face whitish-ochreous. Palpi ochreous-whitish, second joint infuscated. Forewings somewhat dilated, apex obtuse, termen straight, little oblique; 2-5 approximated, 8 to termen; fuscous; basal third brownish-ochreous, edge curved, rather irregular; a suffused pale brownish-ochreous dot on end of cell: cilia fuscous. Hindwings 3 and 4 connate; dark grey; cilia grey.

BRAZIL, Teffé, January (*Parish*); 2 ex.

***Stenoma dilinopa*, n. sp.**

♂ ♀. 20-24 mm. Head, thorax ferruginous-ochreous, face whitish-ochreous. Palpi ochreous tinged ferruginous or fuscous. Forewings suboblong, apex rounded, termen rounded, vertical; 2-5 approximated, 8 to apex (indefinite); pale ochreous; markings formed of suffused brown or ferruginous-brown irroration, viz., a basal patch of which edge runs from $\frac{1}{4}$ of costa to middle of dorsum, separated by a pale streak from a moderate parallel fascia, this nearly followed by an irregular parallel line traversing a linear transverse dark fuscous mark on end of cell, and a præmarginal fascia covering more than $\frac{1}{3}$ of costa and narrowed to tornus, its anterior edge curved; a terminal series of small black dots: cilia pale ochreous, a light brownish antemedian line. Hindwings 3 and 4 connate; light ochreous-yellow, dorsal third more or less tinged grey; cilia pale yellow.

BRAZIL, Teffé, January; PERU, R. Napo, May (*Parish*); 4 ex.

***Stenoma hemiphanta*, n. sp.**

♀. 24-26 mm. Head whitish-ochreous. Palpi whitish, second joint fuscous except towards apex. Thorax pale ochreous. Forewings moderate, somewhat dilated, apex rounded-obtuse, termen slightly rounded, vertical; 2-5 approximated, 8 to apex (indefinite); pale brownish-ochreous, costal edge whitish; a dark fuscous dot on base of costa; three dark fuscous spots on costa, first at $\frac{1}{4}$, small or nearly obsolete, second largest, semioval, beyond

middle, third somewhat smaller, semicircular, at $\frac{4}{5}$; stigmata very small, dark grey, plical slightly beyond first discal, second discal transversely double; a rather curved series of small faint grey dots from third costal spot to tornus; a terminal series of dark fuscous dots: cilia grey with whitish-ochreous reflection. Hindwings 3 and 4 connate; light ochreous-orange, dorsal half grey; cilia whitish-ochreous.

BRAZIL, Teffé, December (*Parish*); 2 ex.

***Stenoma discrepans*, n. sp.**

♀. 22 mm. Head ochreous-whitish, face whiter. Palpi white, second joint suffused light greyish-ochreous except apex. Thorax whitish-ochreous. Forewings apex rounded-obtuse, termen nearly straight, nearly vertical; 2-5 nearly approximated, 8 to just above apex; whitish-yellowish, beyond an irregular line from $\frac{1}{4}$ of dorsum to costa near apex light grey, division suffused; plical and second discal stigmata minute, black; a semioval black spot on middle of costa; a very small black spot on costa at $\frac{4}{5}$, whence a slightly curved series of faint grey dots runs to tornus; a terminal series of small black dots: cilia ochreous-whitish. Hindwings 3 and 4 connate; light grey; cilia ochreous-whitish.

PERU, Jurimaguas, March (*Parish*); 1 ex.

***Stenoma ochrosaris*, n. sp.**

♂ ♀. 17 mm. Head white. Palpi white, second joint with grey lateral streak except towards apex. Thorax light greyish-ochreous, outer margin of patagia whitish. Ovipositor of ♀ broadly flattened laterally. Forewings slightly dilated, costa somewhat sinuate, apex obtuse-rectangular, termen nearly straight, hardly oblique; 2 from near angle, 8 to termen; pale greyish-ochreous, costal edge whitish; three dark fuscous costal spots, first beyond $\frac{1}{4}$, very small, second postmedian, elongate, flattened-triangular, third at $\frac{4}{5}$, somewhat smaller, semioval; stigmata very small, dark fuscous, plical obliquely beyond first discal; an evenly curved series of small indistinct fuscous dots from third costal spot to tornus; an almost marginal series of small dark fuscous dots round apex and termen: cilia whitish-ochreous. Hindwings 3 and 4 connate or closely approximated; ♂ light grey, ♀ grey-whitish; in ♂ a subdorsal groove containing a long expansible yellow-ochreous hairpencil from base; cilia whitish.

BRAZIL, Teffé, January, ♂ (type); Obidos, September, ♀ (*Parish*); probably correctly associated as sexes, and ♀ easily separated from very similar *tricharacta* by ovipositor.

***Stenoma tricharacta*, n. sp.**

♂ ♀. 19-21 mm. Head whitish, second joint light fuscous except apex. Thorax greyish-ochreous. Ovipositor of ♀ slender, cylindrical. Forewings suboblong, hardly dilated, apex obtuse-rectangular, termen nearly straight, vertical; 2-5 approximated, 8 to termen; greyish-ochreous or light fuscous, sometimes tinged

purplish towards dorsum anteriorly, costal edge whitish-ochreous; three semioval or subtriangular dark fuscous costal spots, first at $\frac{1}{4}$, small, others moderate, second median, third about $\frac{4}{5}$; stigmata faintly darker or obsolete, very small; a rather sinuate series of small fuscous dots from third costal spot to tornus; a marginal series of small dark fuscous dots round apex and termen: cilia pale greyish-ochreous, on termen suffused fuscous, with darker median line. Hindwings 3 and 4 connate; grey, in ♀ darker; cilia whitish-grey, a darker subbasal shade.

BRAZIL, Teffé, Obidos, August, September, January; FRENCH GUIANA, R. Maroni; 23 ex. Very like the preceding, but not really very closely related.

Stenoma halmas, n. sp.

♂. 25 mm. Head whitish. Palpi whitish, second joint suffused brownish anteriorly. Thorax light fuscous. Forewings somewhat dilated, apex obtuse, termen nearly straight, hardly oblique; 2-5 approximated, 8 to apex; light fuscous irrorated grey-whitish; extreme costal edge yellowish except towards base; some undefined fuscous suffusion on costa at $\frac{1}{3}$, and semioval dark fuscous spots beyond middle and at $\frac{4}{5}$; a blotch of brownish suffusion on middle of dorsum; second discal stigma dark fuscous; several indistinct darker dots in a subterminal series on lower part of wing; a marginal series of black dots round apex and termen: cilia light yellow-brownish. Hindwings 3 and 4 short-stalked; pale ochreous-yellowish, posterior half suffused grey, terminal edge yellowish; cilia light yellowish, dorsum clothed with pale yellowish hairs.

PERU, Iquitos, March (*Parish*); 1 ex.

Stenoma cathosiota, n. sp.

♀. 31 mm. Head white. Palpi white, basal $\frac{3}{5}$ of second joint suffused grey. Thorax pale lilac-grey. Forewings rather dilated, apex obtuse-rectangular, termen faintly sinuate, hardly oblique; 2-5 approximated, 8 to termen; pale lilac-grey, costal edge ochreous-yellow; stigmata fuscous, plical obliquely beyond first discal; a spot of faint brownish suffusion on middle of dorsum; a small fuscous spot beneath costal edge in middle, whence a series of cloudy undefined fuscous dots rather strongly excurved in disc runs to dorsum about $\frac{3}{4}$; a rather large triangular fuscous spot beneath costal edge towards apex, whence a curved series of small dark fuscous dots runs to tornus; a terminal series of small black dots: cilia grey-whitish, partially suffused lilac-fuscous on termen. Hindwings 3 and 4 connate; pale yellowish, dorsal half tinged very pale grey; cilia yellow-whitish.

BRAZIL, Teffé, January (*Parish*); 1 ex.

Stenoma adoratrix, n. sp.

♂. 25-28 mm. Head white. Palpi white, basal half of second joint faintly tinged brownish. Thorax light violet-brownish becoming white anteriorly. Forewings somewhat dilated, apex

obtuse-pointed, termen sinuate, little oblique; 2-5 nearly approximated at base, 8 to termen; light brownish-ochreous or greyish-ochreous, towards costa and termen suffused whitish, costal edge posteriorly ferruginous-ochreous; an undefined patch of purple-brown suffusion occupying median area of dorsum and extending more than half across wing; a round purplish spot indicating second discal stigma; a rather thick irregular brown streak from middle of costa obliquely outwards to beyond this, and thence rather sinuate to dorsum at $\frac{4}{5}$, and a slender curved streak, tending to be broken into dots from costa at $\frac{3}{4}$ to tornus, discal veins between these streaks sometimes appearing darker-shaded; a terminal series of small indistinct fuscous dots: cilia pale greyish-ochreous, on tornus whitish-tinged, on costa yellowish, on termen a fuscous basal shade. Hindwings 3 and 4 connate or very short-stalked; pale ochreous-yellow, towards dorsum whitish-tinged; cilia whitish-yellow.

BOLIVIA, Prov. del Sara, Dept. Sta. Cruz, 1500 feet, December (*Steinbach*); 1 ex.

Stenoma desidiosa, n. sp.

♂ ♀. 26-27 mm. Head ochreous-whitish. Palpi whitish, basal third of second joint faintly tinged grey. Thorax light violet-brown. Forewings moderately broad, rather dilated, apex obtuse, termen straight, slightly oblique; 2-5 closely approximated, 7 to apex; light brownish, faintly violet-tinged, extreme costal edge whitish; second discal stigma fuscous, tending to be transversely double; undefined cloudy fuscous spots on costa at middle and $\frac{3}{4}$; a marginal series of rather dark fuscous dots round posterior part of costa and termen: cilia whitish-ochreous, a faint greyish shade. Hindwings 3 and 4 connate; pale ochreous-yellowish; cilia whitish-yellowish.

COLOMBIA, R. Tamaua, Choco, 400 feet, February, and R. Negro, 1650 feet, April; 3 ex.

Stenoma negotiosa, n. sp.

♂. 30 mm. Head whitish, centre of crown and lower part of face tinged violet-grey. Palpi whitish, second joint violet-grey except apex. Thorax light greyish-violet, centrally tinged ochreous. Forewings somewhat dilated, apex rounded-obtuse, termen rounded, vertical; 2-5 approximated, 7 to apex (indefinite); light greyish-violet, with slight ochreous tinge; extreme costal edge whitish from $\frac{1}{5}$ to $\frac{2}{3}$; three triangular purplish costal spots, first at $\frac{1}{4}$, very small, second median, small, third about $\frac{4}{5}$, large; stigmata cloudy, purplish, plical obliquely beyond first discal, beneath second discal a small cloudy spot on fold; a rather curved transverse series of faint cloudy purplish dots beyond second discal; an irregularly curved series of faint greyish dots from third costal spot to before tornus; a terminal series of dark fuscous dots: cilia pale grey. Hindwings 3 and 4 connate; grey; cilia whitish-grey.

BRAZIL, Manaos, November (*Parish*); 1 ex.

Stenoma inardescens, n. sp.

♂. 22–23 mm. Head, thorax purplish-grey, face suffused whitish. Palpi whitish, second joint pale greyish-ochreous with streak of violet suffusion. Forewings moderately broad, rather dilated, apex rounded-obtuse, termen straight, hardly oblique; 2 from near angle, 8 to termen; violet-fuscous; a transverse dark brown spot on end of cell, sometimes broken into two, upper part centred brownish-ochreous: cilia light violet-fuscous. Hindwings 3 and 4 connate; dark grey; cilia grey.

BRAZIL, Teffé, January (*Parish*); 2 ex.

Stenoma camptospila, n. sp.

♂. 21–22 mm. Head whitish. Palpi white, second joint light grey except apex. Thorax whitish-grey-ochreous. Forewings dilated, apex obtuse, termen nearly straight, hardly oblique; 2–5 closely approximated, 8 to termen; pale greyish-ochreous; about five small blackish dots in disc in two transverse series at about $\frac{1}{3}$ and before middle; a transverse dark fuscous spot filled with ochreous in disc beyond middle, its lower end produced obliquely towards tornus as a dark fuscous lobe with apex expanded; an undefined cloud of light fuscous suffusion on costa towards apex, whence a hardly traceable faint pale fuscous rather curved shade runs to tornus; a terminal series of small dark fuscous dots: cilia whitish-ochreous, on termen partially suffused pale violet-fuscous. Hindwings 3 and 4 very short-stalked; whitish-yellowish, yellower towards apex; cilia whitish-yellowish.

BRAZIL, Teffé, December, January (*Parish*); 3 ex. Very close to *dispilella*, which is considerably larger (29–30 mm.), with a small dark spot or dot near base, the antemedian dots more numerous and irregular, the postmedian spot larger, broader, wholly transverse without oblique lobe, subterminal line finer, dotted, more distinct, hindwings ochreous-whitish with dorsal third pale grey. Walker's type is without locality, but I have the species from Colombia.

Stenoma satelles, n. sp.

♂. 26 mm. Head, thorax whitish. Palpi white, second joint except apex grey-yellowish with a purplish streak. Forewings somewhat dilated, apex rounded, termen rounded, vertical; 2 from near angle, 8 to apex (indefinite); very pale whitish-grey, costal edge whitish; stigmata grey, plical small, very obliquely beyond first discal, second discal darker; small dark grey spots beneath costal edge at middle and $\frac{4}{5}$, from first a strongly excurved series of indistinct grey dots to dorsum at $\frac{5}{6}$, indented on fold, from second a moderately curved series of grey dots to tornus; a terminal series of dark grey dots: cilia ochreous-whitish. Hindwings 3 and 4 connate; whitish-ochreous, dorsal half very faintly tinged grey; cilia whitish-ochreous.

BRAZIL, Teffé, January (*Parish*); 1 ex.

Stenoma chromolitha, n. sp.

♂. 24 mm. Head, thorax ochreous-white. Palpi whitish, second joint tinged ochreous, with faint greyish streak. Forewings suboblong, apex rounded-obtuse, termen rounded, little oblique; 2-5 approximated, 8 to costa; ochreous-white, costal edge pale greyish-ochreous, towards base dark grey; three dark indigo-blue spots on costa, first two small, at $\frac{1}{4}$ and middle, third larger, at $\frac{4}{5}$; a patch of brownish-grey suffusion covering dorsal half from $\frac{1}{4}$ to tornus; limited posteriorly by a curved dark coppery-grey shade from third costal spot; plical and second discal stigmata forming suffused coppery-purplish spots, each connected with dorsum by a dark coppery-purplish shade, some dark purplish suffusion towards dorsum between these; a marginal series of dark indigo-blue dots round apex and upper part of termen, largest above apex: cilia ochreous-white, round apex suffused fuscous. Hindwings 3 and 4 connate: ochreous-whitish, terminal edge more ochreous-tinged; cilia ochreous-whitish.

BOLIVIA, Prov. del Sara, Dept. Sta. Cruz, 1500 feet; 1 ex.

Stenoma bolistis, n. sp.

♂. 19 mm. Head, thorax white. Palpi white, second joint fuscous except apex. Forewings somewhat dilated, apex rounded-obtuse, termen rounded, vertical; 2-5 approximated, 8 to termen (indefinite); ochreous-white; a small very oblique dark fuscous mark on costa at $\frac{1}{4}$, and a small blackish spot on dorsum at $\frac{1}{4}$, some brownish scales tending to indicate an irregular curved line between these; stigmata very small, indistinct, fuscous, plical obliquely beyond first discal, some faint irregular brownish suffusion towards median area of dorsum; a moderate blackish spot on middle of costa, whence a faint undefined irregular curved line of brownish irroration runs to dorsum at $\frac{3}{4}$; a rather large semioval blackish spot on costa at $\frac{4}{5}$, whence a curved series of dark fuscous dots runs to tornus; a terminal series of dark fuscous sublinear dots: cilia whitish-ochreous. Hindwings 3 and 4 connate; pale whitish-ochreous; cilia concolorous.

BRAZIL, Teffé, December (*Parish*); 1 ex.

Stenoma xylinopa, n. sp.

♂. 28 mm. Head ochreous-whitish, crown faintly brownish-tinged. Palpi white, second joint light grey except apex. Thorax whity-brownish. Forewings suboblong, hardly dilated, apex obtuse, termen straight, nearly vertical; 2 from near angle, 8 to termen; light brown, costal edge obscurely ochreous-whitish; stigmata small, dark fuscous, plical very obliquely beyond first discal; dark fuscous triangular spots on costa beyond middle and at $\frac{4}{5}$, from second a curved series of small indistinct fuscous dots to before tornus; a marginal series of small dark fuscous dots round apex and termen: cilia pale ochreous. Hindwings 3 and 4 connate; dark grey; cilia whitish-grey.

BOLIVIA, Prov. del Sara, Dept. Sta. Cruz, 1500 feet; 1 ex.

Stenoma alluvialis, n. sp.

♂. 24-25 mm. Head white. Palpi white, second joint light fuscous except apex, extreme base of terminal joint dark fuscous. Thorax white sprinkled light purplish-fuscous. Forewings moderate, rather dilated, apex rounded-obtuse, termen rounded, little oblique; 2 from towards angle, 7 to termen; lilac-whitish, costal edge white; dorsal area beneath fold tinged lilac-brownish, with scattered dark fuscous scales, a few lilac-brown scales scattered in disc about stigmata; a fine line of dark fuscous scales along dorsal edge from about $\frac{1}{4}$ to tornus; stigmata dark fuscous, plical very obliquely beyond first discal, second discal transverse-linear; cloudy grey subtriangular spots on costa about middle and $\frac{4}{5}$, from second a curved series of cloudy fuscous dots to tornus, between veins 3-9 a curved series of rather larger similar dots midway between this and end of cell; a marginal series of blackish dots round apex and termen: cilia whitish, base spotted light grey. Hind wings 3 and 4 connate; pale whitish-yellowish; cilia white.

PERU, Papitinga, June (*Parish*); FRENCH GUIANA, R. Maroni; 3 ex.

Stenoma mniodora, n. sp.

♂. 22 mm. Head ochreous-whitish. Palpi whitish, terminal joint with median ring of dark fuscous irroration. Antennæ sinuate near base, ciliations 1. Thorax moss-green. Forewings oblong, apex rounded, termen rounded, nearly vertical; 2-5 approximated, 8 to apex (indefinite): light moss-green; markings deep moss-green; a rather oblique fasciate blotch from dorsum towards base reaching half across wing; a moderate spot on middle of costa; a linear transverse mark on end of cell; a straight transverse subterminal line: cilia whitish-green. Hindwings and cilia green-whitish; 3 and 4 connate.

COLOMBIA, Sosomoko, E. Cordilleras, 2650 feet, March; 1 ex. Next *byssina*.

Stenoma vasifera, n. sp.

♂. 27 mm. Head brownish-ochreous, face whitish-ochreous. Palpi ochreous-whitish, second joint ochreous-greyish. Thorax ochreous-brown. Abdomen dark grey, anal tuft and ventral surface orange. Forewings slightly dilated, costa slightly sinuate in middle, apex obtuse-pointed, termen sinuate, little oblique; 2-5 approximated, 8 to termen; light ochreous-fuscous, becoming more ochreous-tinged posteriorly, a subcostal spot of ochreous suffusion at base; a triangular dark fuscous spot with rounded apex on middle of costa; second discal stigma forming a small transverse-linear fuscous mark; a slight dark brown mark from costa about $\frac{2}{3}$, whence a rather strongly curved series of small indistinct dots runs to near dorsum at $\frac{4}{5}$; dorsal edge tinged brown except towards extremities: cilia concolorous with wing. Hindwings 3 and 4 connate; grey closely irrorated dark grey; cilia whitish-grey, a grey basal shade.

COLOMBIA, Monte del Eden, near Haque, E. Cordilleras, 9600 feet, October; 1 ex. Probably allied to *punicea*.

Stenoma sublimbata Zell.

♂. 28 mm. Head whitish-ochreous. Palpi whitish-ochreous, lower part of second joint infuscated. Antennal ciliations 3. Thorax pale greyish-ochreous mixed brown posteriorly. Forewings moderate, somewhat dilated, apex rounded, termen obliquely rounded; 2-5 approximated, 7 to apex (indefinite); pale yellow-ochreous, with violet gloss, dorsal half and terminal area tinged violet-grey; some narrow violet-grey suffusion along costa except on a patch about $\frac{3}{4}$, and leaving costal edge pale yellowish throughout; a dark brown irregular-edged blotch occupying basal third of dorsum; two small round dark brown spots transversely placed on end of cell, a larger reniform spot beyond upper of these, and a still larger subtriangular spot extending from beyond lower to near dorsum: cilia pale ochreous, suffused light violet-grey on lower part of termen and tornus. Hindwings 3 and 4 connate; ochreous-whitish; a narrow grey marginal streak round posterior $\frac{2}{5}$ of costa and upper part of termen, narrowed and obsolescent downwards; cilia ochreous-grey-whitish, basal half grey round apex and upper part of termen.

COLOMBIA, Monte del Eden, 9600 feet, October; 1 ex. This and the next species are allied to *castellana*. I am indebted to the great generosity of Prince Caradja for the types of these and a large number of other interesting species from Colombia, Bolivia, and Peru.

Stenoma ancylacma, n. sp.

♀. 24 mm. Head, thorax dark fuscous, face whitish. Palpi fuscous. Forewings somewhat dilated, apex rounded, termen rather obliquely rounded; 2-5 approximated, 7 to costa; white, dorsal half suffused pale ochreous and sprinkled fuscous posteriorly, costal edge tinged pale ochreous; a blackish blotch occupying basal third of dorsum and extending gradually and irregularly narrower to base of costa; plical stigma blackish; a triangular brown patch resting on termen from near apex to tornus and its apex reaching in disc to end of cell, upper part yellow-brown and edged posteriorly with some dark fuscous scales and anteriorly with a semi-oval blackish spot, preceding its apex a dark fuscous dot in disc; 3 or 4 indistinct dots of dark fuscous irroration just beneath costa towards apex: cilia whitish, a fuscous patch in middle of termen, beneath this suffused light brownish. Hindwings 3 and 4 connate; ochreous-whitish; a short dark grey almost marginal streak round apex; cilia whitish, round apex base dark grey.

PERU, Andes, at high elevation; 1 ex.

Stenoma orthographa, n. sp.

♂. 24 mm. Head, palpi pale fuscous. Thorax fuscous. Forewings suboblong, slightly dilated, apex rounded, termen rounded, vertical; 2 from near angle, 8 to apex (indefinite); fuscous; three cloudy dark fuscous lines, first two nearly straight, first scarcely oblique, second rather oblique, third forming a series of cloudy dots,

somewhat curved, indented above middle: cilia light fuscous. Hindwings 3 and 4 short-stalked; grey; cilia light grey.

BRAZIL, Orientes (Upper Amazon), February (*Parish*); 1 ex.

***Stenoma pleurotricha*, n. sp.**

♂. 29–30 mm. Head whitish-fleshcolour. Palpi ochreous-whitish, second joint fuscous except towards apex. Thorax light greyish-fleshcolour. Forewings slightly dilated, apex rounded-obtuse, termen slightly rounded, hardly oblique; 2 from near angle, 8 to apex (indefinite); light greyish-fleshcolour; costal edge whitish; plical and second discal stigmata dark grey; a curved subterminal series of cloudy grey dots, sinuate inwards above middle; a marginal series of dark grey dots round apex and termen, surrounded with whitish tinge: cilia whitish-ochreous. Hindwings 3 and 4 connate; grey; a fringe of long pale ochreous-yellowish expansible hairs from near dorsal edge on basal half projecting directly inwards beneath abdomen; cilia whitish-grey.

BRAZIL, Teffé, January (*Parish*); 3 ex.

***Stenoma augescens*, n. sp.**

♀. 39 mm. Head light brownish-ochreous, face whitish-tinged. Palpi whitish, second joint fuscous except apex. Thorax light brown. Forewings somewhat dilated, apex rounded-obtuse, termen vertical; 2 from near angle, 7 to termen, 8 to costa; light brown; costal edge pale ochreous; plical stigma very small, elongate, dark fuscous, second discal forming a rather large roundish dark fuscous spot; a subterminal series of indistinct fuscous subrescentic dots sinuate inwards towards costa; a marginal series of dark fuscous dots round apex and termen: cilia pale brownish-ochreous. Hindwings 3 and 4 connate; rather dark grey; cilia pale grey, a darker subbasal shade.

PERU, Putomayo, R. Ampiyacu; 1 ex.

***Stenoma deltomis*, n. sp.**

♂. 33 mm. Head whitish. Palpi whitish, second joint dark grey except apex. Thorax whitish-grey. Forewings suboblong, slightly dilated, costa on basal third with triangular raised or partially reflexed flap fringed with rough scales towards its apex, then straight, apex rectangular, termen faintly sinuate, vertical; 2 from near angle, 7 to termen, 8 to costa; iridescent-whitish-grey; costal edge ochreous-white from flap; stigmata grey, plical obliquely beyond first discal; a strongly curved series of cloudy greyish dots from towards costa at $\frac{3}{5}$ to near dorsum at $\frac{3}{4}$, interrupted in middle by a round cloudy light grey-brownish spot; a strongly curved subterminal series of fuscous dots, sinuate inwards towards costa; a terminal series of blackish-grey dots: cilia whitish. Hindwings 3 and 4 very short-stalked; light grey, becoming light greyish-ochreous towards apex; cilia ochreous-whitish.

BRAZIL, Teffé and Santa Rita, January, February (*Parish*); 2 ex.

Stenoma incitatrix, n. sp.

♂. 28 mm. Head whitish-ochreous. Palpi whitish, basal $\frac{2}{3}$ of second joint fuscous. Thorax pale pinkish-ochreous. Forewings somewhat dilated, apex rounded-obtuse, termen hardly oblique; 2 from near angle, 7 to termen, 8 to costa; pale pinkish-ochreous; costal edge ochreous-whitish; plical and second discal stigmata dark grey; a marginal series of dark grey dots round apex and termen; cilia ochreous-whitish. Hindwings 3 and 4 short-stalked; ochreous-whitish, tinged greyish-ochreous towards apex; a fringe of long whitish-ochreous expansible hairs from near dorsum on upper half projecting inwards beneath abdomen; cilia ochreous-whitish, a faint greyish subbasal shade.

ARGENTINA, R. Parana; 1 ex. Allied to *pleurotricha*.

Stenoma relata, n. sp.

♂ ♀. 19–21 mm. Head whitish, crown sometimes tinged grey. Palpi white, second joint fuscous except apex, base and subapical ring of terminal joint dark fuscous. Thorax whitish-grey. Forewings slightly dilated, apex rounded-obtuse, termen rounded, nearly vertical; 2 from near angle, 7 to termen, 8 to costa; whitish-grey; costal edge white; stigmata blackish-grey; plical obliquely beyond first discal; some faint grey speckling indicating an oblique shade about $\frac{2}{5}$; an irregular slightly curved grey shade from middle of costa behind second discal stigma to dorsum at $\frac{4}{5}$; a curved series of dark grey dots from costa at $\frac{3}{4}$ to dorsum before tornus, strongly indented towards costa; a marginal series of blackish dots round apex and termen; cilia white, tinged ochreous towards base. Hindwings 3 and 4 very short-stalked; pale grey; cilia whitish, a greyish subbasal line.

BRAZIL, Teffé, January; FRENCH GUIANA, R. Maroni; PERU, Papitinga (Upper Amazon), January (*Parish*); 5 ex.

Stenoma perjura, n. sp.

♂. 21 mm. Head ochreous-whitish. Palpi white, basal $\frac{2}{3}$ of second joint fuscous. Thorax whitish-grey-ochreous. Forewings suboblong, hardly dilated, apex rounded, termen vertical; 2 from near angle, 7 to termen, 8 to costa; whitish-grey-ochreous; costal edge ochreous-whitish; stigmata dark fuscous, plical obliquely beyond first discal but unusually near it; faint fuscous suffusion indicating an irregular shade from beyond second discal stigma to dorsum; a curved subterminal series of indistinct cloudy fuscous dots sinuate-indented above middle; a terminal series of dark fuscous dots; cilia whitish. Hindwings 3 and 4 connate; pale grey, whitish-suffused towards base; a pale greyish-ochreous subcostal expansible hairpencil from base to near middle (but costal area not expanded); cilia ochreous-whitish, a grey subbasal shade round apex and upper part of termen.

BRAZIL, Teffé, January (*Parish*); 1 ex.

Stenoma salubris, n. sp.

♂. 13-14 mm. Head white. Palpi white, second joint grey except apex, base of terminal joint dark fuscous. Thorax grey-whitish. Forewings slightly dilated, apex rounded-obtuse, termen rounded, somewhat oblique; 2-5 approximated, 7 to apex; grey-whitish; a blackish dot on base of fold; stigmata rather large, blackish, plical obliquely beyond first discal; some slight fuscous suffusion towards base of dorsum; a cloudy dark fuscous shade connecting plical stigma with dorsum; a suffused dark fuscous shade from costa at $\frac{3}{5}$ somewhat angulated beyond second discal stigma to dorsum near tornus; a curved dark fuscous line from costa at $\frac{4}{5}$ to tornus, indented towards costa; a marginal series of blackish dots round apex and termen: cilia white. Hindwings 3 and 4 connate; pale grey; cilia whitish.

BRAZIL, Parintins, Teffé, October, December (*Parish*); 4 ex.

Stenoma machinatrix, n. sp.

♂. 28 mm. Head, thorax pale brownish-ochreous, face whitish-tinged, thoracic crest tinged ferruginous-brown on sides. Palpi ochreous-whitish. Forewings suboblong, costa anteriorly moderately arched, apex rounded, termen rounded, nearly vertical; 2 from near angle, 7 to apex (indefinite); pale brownish-ochreous; dorsal subbasal projecting scales tinged ferruginous-brown; an oblique series of four irregular cloudy dark fuscous dots from costa at $\frac{1}{4}$ to a small cloudy ferruginous-brown spot on fold, thence continued by a faint sinuate fuscous shade to dorsum at $\frac{1}{3}$; a straight ferruginous shade suffused posteriorly from before middle of costa to tornus, two obliquely placed cloudy darker dots on anterior edge in middle; an irregular waved faint fuscous line from $\frac{2}{3}$ of costa to tornus, where it meets preceding, excurved in middle, on lower portion with two or three crescentic dark fuscous marks; a terminal series of dark fuscous dots: cilia pale brownish-ochreous. Hindwings 3 and 4 connate; pale grey; cilia whitish-ochreous.

COLOMBIA, Minero, 1650 feet, October; 1 ex.

Stenoma finitrix, n. sp.

♂. 18-21 mm. Head whitish, back of crown grey. Palpi grey, apex of second joint whitish, terminal joint whitish towards base and posteriorly. Thorax dark purplish-grey. Forewings sub-oblong, apex rounded, termen rounded, little oblique; 2 from near angle, connected with 1c, 7 to costa; dark purplish-grey; costa slenderly pale ochreous-yellow from base to $\frac{4}{5}$, whence a marginal series of white marks extends round apex and termen; three cloudy whitish dots in an indented series from middle of costa, a crescentic mark on end of cell, and a cloudy dot on fold beneath this: cilia light grey, outer half white on termen, becoming yellowish on costa. Hindwings 3 and 4 moderately stalked; dark grey; cilia dark grey, a darker subbasal shade.

COLOMBIA, San Antonio, 6600 feet, February; BOLIVIA, R. Songo, 2800 feet; 3 ex.

Stenoma horocyma, n. sp.

♂. 10 mm. Head white. Palpi white, basal half of second joint, and base and subapical ring of terminal joint dark fuscous. Antennal ciliations 4. Thorax dark indigo-grey, anterior margin white. Forewings suboblong, somewhat narrowed posteriorly, apex rounded, termen rounded, vertical; 7 to apex (indefinite); glossy dark indigo-blue-grey; costal edge white; white marks on costa before and beyond middle; second discal stigma white; a waved white line just within margin round posterior part of costa and termen: cilia white, basal third grey. Hindwings 4 absent, 3 and 5 connate; grey; cilia whitish, a grey subbasal shade.

BRAZIL, Teffé, December (*Parish*); 1 ex.

Stenoma icteropis, n. sp.

♂ ♀. 13-14 mm. Head pale ochreous somewhat mixed blue-grey. Palpi whitish-ochreous, terminal joint more or less grey anteriorly. Thorax grey, shoulders pale ochreous. Forewings suboblong, apex rounded, termen rounded, little oblique; 2 from near angle, 7 to costa (indefinite); pale violet-grey, sometimes suffused whitish-ochreous; a short yellow-ochreous mark from middle of base; a slender yellow-ochreous costal streak from base to about middle; a suffused ochreous-whitish bar on end of cell: cilia whitish-ochreous, yellower towards base. Hindwings 4 absent, 3 and 5 approximated or connate; light grey; cilia whitish, a greyish subbasal line.

BRAZIL, Santarém, August (*Parish*); 3 ex.

Stenoma vitreola, n. sp.

♂. 12 mm. Head, thorax pale silvery-grey, face silvery-white. Palpi white, second joint with grey streak to near apex, terminal joint greyish on apical half, base dark grey. Forewings rather elongate, apex rounded, termen rather obliquely rounded; 3 and 4 out of 2, 7 to apex (indefinite); light silvery-grey, with whitish reflections between veins posteriorly, dorsal area darker grey; costal edge white; second discal stigma cloudy, dark grey, from beneath it an oblique white streak to near dorsum before tornus: cilia ochreous-whitish, an interrupted grey subbasal shade. Hindwings 3 and 4 connate, 5 nearly parallel; light grey; cilia grey-whitish, a greyish subbasal shade.

BRAZIL, Para, July (*Parish*); 1 ex. I have also from Obidos 2 ♀ specimens, which are probably the other sex of this species; 13 mm., forewings slightly broader posteriorly, more brownish-tinged, second discal stigma black, distinct, white streak less developed but edged anteriorly by a triangular dorsal blotch of dark fuscous suffusion. Allied to *stylonota*.

Stenoma atmospora, n. sp.

♀. 38-39 mm. Head whitish-ochreous. Palpi whitish-ochreous tinged brownish. Thorax pale greyish-ochreous. Forewings

elongate-oblong, apex rounded-obtuse, termen slightly oblique; 2-5 approximated, 7 to apex; pale greyish-ochreous sprinkled dark brown specks, these forming a darker cloud towards median area of dorsum; second discal stigma rather large, dark fuscous; terminal dots sometimes slightly indicated: cilia pale ochreous. Hindwings 3 and 4 short-stalked; pale grey; cilia pale greyish-ochreous.

COLOMBIA, Minero and Sosomoko, 2650 feet, April, October; 2 ex.

Stenoma recurrens, n. sp.

♂ ♀. 19-20 mm. Head pale ochreous, face suffused whitish. Palpi whitish-ochreous, a grey streak on basal half of second joint. Thorax pale brownish-ochreous. Forewings oblong, apex rounded-obtuse, termen vertical; 2-5 approximated, 7 to termen, 8 to costa; light brownish or brownish-ochreous; plical and second discal stigma small, dark fuscous; three very irregular waved-dentate slender lines of dark fuscous irroration, first slightly curved from costa at $\frac{1}{3}$ to dorsum at $\frac{2}{5}$, second suffused brownish, from middle of costa obtusely angulated just beyond second discal stigma to dorsum at $\frac{2}{3}$, third forming a series of dark dots connected by a waved line towards costa, from costa at $\frac{2}{3}$ strongly excurved to dorsum near tornus, indented above middle and sinuate on lower portion; a marginal series of blackish dots round posterior part of costa and termen, on apical part of costa surrounded whitish suffusion: cilia ochreous-grey, sometimes obscurely barred whitish, on apex and costa suffused whitish. Hindwings 3 and 4 connate; grey; cilia whitish.

BOLIVIA, Prov. del Sara, Dept. Santa Cruz, 1500 feet; 4 ex.

Stenoma agraria, n. sp.

♂ ♀. 20-22 mm. Head, thorax light fuscous, face whitish-tinged. Palpi whitish-fuscous. Forewings oblong, apex rounded-obtuse, termen vertical; 2-5 approximated, 7 to termen, 8 to costa; light fuscous; plical and second discal stigmata small, cloudy, dark fuscous; three cloudy darker transverse curved shades, first from $\frac{1}{3}$ of costa to $\frac{2}{5}$ of dorsum, second from middle of costa to $\frac{2}{3}$ of dorsum, third from $\frac{3}{4}$ of costa to near tornus; a terminal series of small dark fuscous dots: cilia greyish-ochreous. Hindwings 3 and 4 connate; grey; cilia grey-whitish.

BOLIVIA, Prov. del Sara, Dept. Santa Cruz, 1500 feet; 2 ex.

Stenoma melixesta, n. sp.

♀. 42 mm. Head, thorax pale ochreous, face suffused whitish. Palpi whitish-ochreous, second joint infuscated except towards apex. Forewings suboblong, slightly dilated, apex obtuse, termen vertical; 2 from near angle, 8 to termen; ochreous, somewhat lighter towards costa anteriorly; dorsal edge fuscous from near base to termen; plical and second discal stigmata small, dark fuscous, first discal indicated by faint fuscous suffusion; a faintly indicated streak of slight fuscous suffusion from middle of costa

to dorsum at $\frac{2}{3}$, angulated beyond second discal stigma; a gently curved line of fuscous scales from costa at $\frac{3}{4}$ to dorsum towards tornus: some slight fuscous speckling towards termen: cilia light grey tinged ochreous, tips pale. Hindwings 3 and 4 connate; pale grey, apical area suffused whitish-yellowish; cilia pale yellowish becoming whitish towards tips.

COLOMBIA, Rio Negro, 1650 feet, April; 1 ex.

Stenoma ischioptila, n. sp.

♂. 30 mm. Head, thorax light fuscous, face whitish. Palpi whitish, externally suffused fuscous. Forewings somewhat dilated, apex rounded-obtuse, termen rounded, little oblique; rather light fuscous; extreme costal edge ochreous-white; plical and second discal stigmata blackish; a faintly indicated hardly curved darker shade from middle of costa just beyond second discal stigma to dorsum at $\frac{2}{3}$; a curved subterminal series of faint cloudy greyish dots sinuate towards costa; a marginal series of blackish dots round apex and termen: cilia whitish-fuscous. Hindwings 3 and 4 connate; dark grey; an expansible whitish subcostal hairpencil from base to middle (costal area not expanded); cilia light grey. Abdomen with dorsal tuft of long grey hairs towards base, and a large expansible pencil of long ochreous-whitish hairs from each side at base lying in a subdorsal groove of hindwings surrounded with long grey hairs.

COLOMBIA, Sosomoko, 2650 feet, February; 1 ex.

Stenoma trymalopa, n. sp.

♂. 30 mm. Head, thorax light fuscous, face suffused whitish. Palpi whitish, a grey streak on second joint to $\frac{2}{3}$. Forewings suboblong, slightly dilated, apex rounded-obtuse, termen vertical; 2 from near angle, 7 to termen, 8 to costa; light fuscous; discal stigmata small, dark grey; three cloudy rather dark grey lines from beneath costa, first oblique to first discal stigma, second oblique to second discal, third tending to be interrupted into dots, curved-oblique to tornus, somewhat indented above middle and irregular near tornus; a pale yellowish rather oblique-reniform spot resting on middle of dorsum; a marginal series of dark fuscous dots round apex and termen: cilia greyish. Hindwings 3 and 4 connate; grey; cilia pale greyish.

COLOMBIA, San Antonio, 6600 feet, February; 1 ex.

Stenoma germinans, n. sp.

♂. 27 mm. Head, thorax pale brownish, face whitish-tinged. Palpi whitish, second joint infuscated to above middle. Forewings slightly dilated, apex rounded-obtuse, termen vertical; 2-5 approximated, 7 to termen, 8 to costa; light brownish-fuscous; well-marked scaletufts above and below middle at $\frac{1}{5}$; plical and second discal stigmata dark fuscous; fasciæ of very faint darker suffusion crossing wing at $\frac{1}{3}$ and beyond middle, second slightly curved; a rather curved waved fuscous line from costa at $\frac{4}{5}$ to dorsum

before tornus, sinuate towards costa: cilia pale ochreous, tips whitish. Hindwings 3 and 4 connate; grey, darker posteriorly; an expansible fringe-tuft of long pale ochreous hairs from near dorsum on upper half projecting inwards beneath abdomen; cilia grey-whitish, a grey subbasal shade.

COLOMBIA, locality uncertain; 1 ex. The tufts of forewings are unique, but the normal palpi and other characters show that it is not a *Ptilogenes*. Allied to *pleurotricha*.

Stenoma iostalacta, n. sp.

♂. 24 mm. Head ochreous-whitish. Palpi whitish, second joint greyish-ochreous except apex. Antennal ciliations 2. Thorax whitish-ochreous. Forewings somewhat dilated, apex rounded, termen vertical; 2 from near angle, 7 to termen, 8 to costa; pale grey; costal edge pale ochreous; markings light ferruginous-brownish, darker towards costa and partially on edges; a dot near base in middle; a triangular patch extending on dorsum from near base to middle, its apex confluent with a spot on costa at $\frac{1}{3}$, its posterior edge with a projection in disc; a transverse-oval spot representing second discal stigma; a triangular spot on middle of costa, whence a shade strongly excurved beyond second discal stigma runs to dorsum at $\frac{3}{4}$, its extremity produced each way on dorsal edge; a larger triangular spot on costa at $\frac{3}{4}$, whence a gently curved series of small cloudy spots runs to tornus; a marginal series of blackish dots round apex and termen: cilia pale brownish-ochreous becoming whitish towards tips. Hindwings 3 and 4 connate; light grey; cilia grey-whitish, a pale greyish subbasal shade.

PERU, Cocapata, 12,000 feet; 1 ex. Close to *compressa*.

Stenoma chromotechna, n. sp.

♂. 17 mm. Head, thorax pale greyish-ochreous, face whitish, patagia dark brown. Palpi white, second joint dark grey except apex, terminal joint grey anteriorly towards apex, base dark grey. Forewings suboblong, costa hardly sinuate, apex rounded, termen nearly vertical; 2-5 approximated, 7 to apex; light greyish-violet; basal third of dorsum and projecting scales ferruginous-brown; a fine white longitudinal line from base just above this extending to middle; costal edge whitish; a slender blackish streak from costa at $\frac{1}{3}$ to dorsum near tornus, expanded on costa, partially tinged brownish on its edges, margined with a whitish streak anteriorly to fold and posteriorly beyond fold, preceded on fold by a small greyish and ochreous tuft; space between this and next streak mixed or tinged whitish-ochreous; an oblique dark brown white-margined fasciate streak from middle of costa reaching half across wing, then becoming black and continued by a blackish line to dorsum where it almost meets apex of preceding; apical fourth of wing forming a blotch with anterior edge slightly convex, its anterior half whitish, posterior pale ochreous; a marginal series of blackish dots round apex and termen: cilia pale ochreous, a faint

brownish line. Hindwings 3 and 4 short-stalked; rather dark grey; cilia grey-whitish, a grey subbasal shade.

BRAZIL, Teffé, January (*Parish*); 1 ex. A singular and elegant form, perhaps allied to *modulata*.

***Stenoma sciocnesta*, n. sp.**

♂ ♀. 13–14 mm. Head pale ochreous-grey, face glossy, whitish-tinged. Palpi ochreous-whitish suffusedly mixed grey. Thorax light grey. Forewings rather elongate, apex rounded-obtuse, termen rounded, somewhat oblique; 2–5 approximated, 7 to apex (indefinite); pale ochreous-grey irregularly sprinkled dark grey; stigmata dark fuscous, plical obliquely beyond first discal; cloudy spots of grey suffusion on costa at $\frac{1}{3}$, beyond middle, and $\frac{4}{5}$, from third a strongly curved series of cloudy dark grey subconfluent dots to tornus: cilia light greyish, a dark grey subbasal line. Hindwings 3 and 4 rather short-stalked; grey; cilia light grey, a darker subbasal shade.

BRAZIL, Teffé, December, January (*Parish*); 3 ex. Close to *epicnesta* (which also occurs at Teffé), but easily known by the grey (not whitish) groundcolour, and absence of terminal dots.

***Stenoma tyroxesta*, n. sp.**

♂. 14 mm. Head grey, face glossy ochreous-whitish. Palpi grey. Thorax pale glossy greyish-ochreous. Forewings hardly dilated, apex rounded, termen vertical; 2–5 approximated, 7 to apex (indefinite); pale glossy greyish-ochreous; costa suffused grey on basal fourth; a dark grey dot near base in middle; stigmata small, dark grey, plical obliquely beyond first discal; an oblique grey mark on middle of costa; a rather strongly curved series of blackish dots from costa at $\frac{3}{4}$ to tornus, slightly sinuate towards costa and below middle; a marginal series of blackish dots round apex and termen: cilia pale greyish-ochreous, a dark grey subbasal line. Hindwings 3 and 4 short-stalked; grey; cilia light grey.

BRAZIL, Teffé, January (*Parish*); 1 ex.

***Stenoma congrua*, n. sp.**

♂. 15 mm. Head white. Palpi white, second joint fuscous except apex, base of terminal joint dark fuscous. Thorax ochreous-white. Forewings rather elongate, apex obtuse, termen straight, little oblique; 2–5 approximated, 7 to apex; ochreous-white, some irregular fuscous sprinkling, especially beyond cell; a slight dark fuscous mark on base of costa; second discal stigma blackish; three oblique blackish costal spots, first at $\frac{1}{3}$, small, second in middle, moderate, third at $\frac{4}{5}$, elongate-semioval, from end of this a somewhat curved dentate fuscous shade near termen to tornus; a marginal series of black dots round apex and termen: cilia whitish, a light fuscous subbasal line. Hindwings 3 and 4 short-stalked; light grey, terminal edge suffused darker; cilia whitish, a grey subbasal line.

PERU, Iquitos, May (*Parish*); 2 ex.

Stenoma associata, n. sp.

♂. 16–17 mm. Head white, crown tinged grey. Palpi white somewhat sprinkled dark fuscous, second joint with grey streak on basal half and subapical ring, terminal joint with blackish anterior and interior lines. Thorax whitish-grey. Forewings rather elongate-oblong, apex rounded-obtuse, termen vertical; 2–5 approximated, 6 and 7 slightly approximated at tip, 7 to apex (indefinite); ochreous-grey-whitish; second discal stigma dark fuscous; sometimes some fuscous irroration towards base of costa and dorsum; an oblique blackish-grey mark on costa at $\frac{1}{3}$, whence a faint interrupted brownish shade runs to some suffusion on dorsum about middle; a triangular blackish-grey spot on middle of costa, whence a similar faint shade runs to some dorsal suffusion at $\frac{3}{4}$; a larger triangular blackish-grey spot on costa about $\frac{3}{4}$, whence a curved brownish line runs to tornus; some black marginal dots round apex and termen, largest at and above apex, on termen connected by a fuscous line: cilia whitish or grey-whitish, a dark fuscous subbasal shade at and above apex. Hindwings 3 and 4 short-stalked; light greyish; cilia whitish.

BRAZIL, Para, Santarem, August; PERU, Jurimaguas, March (*Parish*); 4 ex.

Stenoma paraplecta, n. sp.

♂. 13 mm. Head, palpi, thorax grey, face silvery-metallic. Forewings rather elongate, apex rounded, termen rounded, somewhat oblique; 2–5 approximated, 7 to costa; violet-grey, costa and dorsum suffused darker, from base to end of cell mostly suffused obscurely whitish-ochreous except on costa and dorsum, and more vaguely to about $\frac{4}{5}$, between cell and posterior half of costa with four pale ochreous-yellowish gradually expanded interneural streaks ending abruptly before reaching costa; a fine black marginal line round apex and termen preceded by slender silvery-whitish-ochreous suffusion: cilia pale yellowish, outer half pale silvery-grey. Hindwings costa somewhat sinuate before abrupt origin of cilia, 3 and 4 moderately stalked; grey; a dark grey marginal line round apex; cilia grey, round apex basal half whitish-yellowish.

BRAZIL, Teffé, January (*Parish*); 1 ex. An eccentric species.

Stenoma secundata, n. sp.

♂. 12 mm. Head white. Palpi white, second joint dark fuscous except apex. Thorax fuscous-whitish. Forewings hardly dilated, apex rounded-obtuse, termen slightly oblique; 2–5 approximated, 7 to apex (indefinite); whity-brownish; second discal stigma dark fuscous, plical indicated by one or two scales; small triangular dark fuscous spots on costa at middle and $\frac{3}{4}$, from second a curved series of faint dots of fuscous irroration to tornus: a marginal series of black dots round apex and termen: cilia ochreous-whitish, a faint fuscous median line and apical spot. Hindwings 3 and 4 connate; grey; cilia pale grey, a darker subbasal shade.

PERU, Jurimaguas, March (*Parish*); 1 ex.

Stenoma hexascia, n. sp.

♂. 15 mm. Head, thorax shining pearl-white. Palpi whitish, towards base slightly infuscated. Forewings slightly dilated, apex obtuse, termen rounded, vertical; 2-5 approximated, 7 to termen, 8 to costa; white; five small suffused blackish-grey costal spots, first slightly marked, basal, fifth at $\frac{4}{5}$, first three and fifth giving rise to cloudy suffused greyish-ochreous transverse fasciæ, fifth running to tornus, fourth to two similar fasciæ confluent beneath costa with third and fifth also, between these two second discal stigma transverse, blackish, plical indicated by one or two blackish scales; beyond fifth two blackish costal dots, round apex and upper part of termen a greyish-bronzy line: cilia greyish-ochreous, towards tips whitish. Hindwings 3 and 4 short-stalked; pale greyish; cilia grey-whitish, a light grey subbasal line.

BRAZIL, Teffé, December, January (*Parish*); 2 ex.

Stenoma antilyra, n. sp.

♂ ♀. 15-17 mm. Head, thorax white. Palpi white, a fuscous streak on basal half of second joint. Forewings gradually dilated, apex obtuse, termen straight, slightly oblique; 2 and 3 stalked, 7 to termen, 8 to costa; white, variably tinged or sprinkled greyish, in one specimen obscurely streaked pale greyish-ochreous between veins; costal edge dark grey towards base; some irregular greyish clouding towards costa at $\frac{1}{4}$; a very oblique dark grey mark from before middle of costa, beyond this two small grey spots edged darker, from beyond second a series of dark grey dots obliquely inwards, forming a deep indentation, then strongly ex-curved in disc and again sinuate to before tornus; first discal stigma indistinct, fuscous, plical obliquely beyond it, indistinct or seldom dark fuscous, second discal represented by two transversely placed dark fuscous dots, an additional dot before lower of these; two dark fuscous marks on costa before apex, and a terminal series of dots or interrupted line: cilia grey, tips whitish. Hindwings 3 and 4 connate; whitish or whitish-grey; a dark fuscous terminal line; cilia grey-whitish.

BRAZIL, Parintins, Manaus, Teffé, October to January; PERU, R. Napo, May (*Parish*); FRENCH GUIANA, R. Maroni; 6 ex.

Stenoma robiginosa, n. sp.

♂. 16 mm. Head white, face tinged pink. Palpi white, second joint pinkish-grey, base and subapical ring of terminal joint dark fuscous. Thorax greyish-fleshcolour, shoulders suffused white. Forewings rather dilated, apex rounded-obtuse, termen vertical; 2-5 approximated, 7 to termen, 8 to costa; pale grey; costal edge ochreous-white, with oblique dark grey marks at base, $\frac{1}{4}$, and $\frac{1}{3}$, a spot in middle, and larger triangular spot at $\frac{4}{5}$; an irregularly triangular light pinkish-brown patch extending on dorsum from base to middle, its apex reaching nearly to median costal spot; second discal stigma dark fuscous; an irregular light pinkish-brown shade from median costal spot obtusely angulated in middle

to dorsum at $\frac{3}{4}$, and a curved shade from posterior costal spot to tornus: a marginal series of blackish dots round apex and termen: cilia light brownish-grey. Hindwings 3 and 4 connate; grey; cilia grey-whitish.

BRAZIL, Manaos, November (*Parish*); 1 ex.

***Stenoma chloroxantha*, n. sp.**

♀. 21 mm. Head pale ochreous. Palpi ochreous-whitish, second joint fuscous except apex. Thorax ochreous-brown. Forewings suboblong, termen slightly sinuate, little oblique; 2 from towards angle, 7 to termen, 8 to costa; light ochreous-brown, with faint pinkish tinge; plical stigma forming a round pale green spot edged fuscous, second discal an indistinct pale green dot; a light yellow triangular terminal patch, extending on dorsum to $\frac{2}{3}$ and narrowed to a point at apex, edged slight fuscous suffusion anteriorly, marked with a fine light yellow-brown slightly curved line from anterior edge above middle to dorsum before tornus, and a terminal series of brownish dots: cilia yellow-ochreous, with faint basal greyish line, on costa greyish-suffused, on tornus whitish-ochreous. Hindwings 3 and 4 connate; pale grey; cilia grey-whitish, a faint greyish subbasal line.

BRAZIL, Rio Grande do Sul, Santa Cruz; 1 ex.

***Stenoma hoplitica*, n. sp.**

♂. 16-17 mm. Head, thorax light grey-pinkish, face white. Palpi white, tinged pinkish-grey except apex of joints. Forewings somewhat dilated, apex obtuse, termen straight, slightly oblique; 2 from near angle, 8 to apex; light pinkish-violet; costal edge ochreous-yellow to near apex; plical stigma obscurely grey, followed by some whitish scales, second discal transverse, dark grey; a suffused pale grey terminal patch, its edge rather convex, running from apex to dorsum about $\frac{2}{3}$, traversed by a grey sinuate subterminal shade: cilia light purple-grey, towards tornus whitish-tinged. Hindwings 3 and 4 connate; grey; cilia whitish-grey.

BRAZIL, Teffé, December, January (*Parish*); 2 ex.

***Stenoma dryaula*, n. sp.**

♀. 14-15 mm. Head ochreous-whitish. Palpi whitish, a dark fuscous streak on second joint to near apex. Thorax whitish-ochreous. Forewings elongate, apex obtuse, termen rather obliquely rounded; 2 and 3 stalked, 7 to apex; whitish-ochreous, dorsal half slightly brownish-tinged; a moderate brown median streak from base to apex, upper edge somewhat mixed dark fuscous, lower rather suffused; second discal stigma black, plical also but smaller and sometimes obsolete; a marginal series of black dots round apical part of costa and termen: cilia whitish-ochreous, at apex a brownish bar. Hindwings 3 and 4 short-stalked; light grey; cilia whitish-grey.

BRAZIL, Obidos, August, September (*Parish*); 12 ex., all ♀.

Stenoma procritica, n. sp.

♂ ♀. 22–29 mm. Head white, crown tinged brownish. Palpi white, second joint light brownish except apex. Thorax white more or less suffused brownish. Forewings rather elongate, apex rounded-obtuse, termen rounded, slightly oblique; 2 from towards angle, 7 to termen, 8 to costa; white, variably suffused pale brownish, which forms slender costal and dorsal streaks and moderately broad undefined longitudinal streaks above and below middle throughout, these cut by pale or white lines on veins and more or less mixed blackish-fuscous between them; stigmata black, plical obliquely beyond first discal, a fine blackish line along fold from base to this; a curved subterminal series of cloudy dots of fuscous or dark fuscous irroration; a marginal series of black dots round apex and termen: cilia ochreous-whitish, faintly barred pale ochreous. Hindwings 3 and 4 connate; pale greyish, whitish-tinged towards base, apical edge suffused dark grey; cilia grey-whitish, a grey subbasal line.

BRAZIL, Teffé, December (*Parish*); S.E. PERU, Santo Domingo, 6000 feet, November (*Ockenden*); 2 ex.

Stenoma conveniens, n. sp.

♀. 14 mm. Head, thorax pale ochreous, face glossy whitish-ochreous. Palpi ochreous-whitish, second joint with fuscous streak to $\frac{3}{4}$, base of terminal joint fuscous. Forewings elongate, apex rounded, termen rounded, little oblique; 2 and 3 connate, 7 to apex (indefinite); whitish-ochreous, all veins marked brownish-ochreous lines, these not reaching margin but connected by a curved præ-terminal shade, on lower angle of cell a slight thickening of dark fuscous suffusion; a marginal series of black dots round apical part of costa and termen: cilia whitish-ochreous. Hindwings 3 and 4 moderately stalked; light grey; cilia whitish-grey.

BRAZIL, Matura, June (*Parish*); 1 ex.

Stenoma leptogma, n. sp.

♂. 29 mm. Head, thorax whitish-ochreous. (Palpi broken.) Forewings elongate, apex almost rectangular, termen slightly rounded, somewhat oblique; 2 from near angle, 7 to termen, 8 to costa; whitish-ochreous, veins marked very fine light brown lines, fainter brownish-ochreous interneural lines between these; very small dark grey dots on both angles of cell: cilia ochreous-whitish. Hindwings 3 and 4 very short-stalked; pale whitish-grey; cilia whitish.

COLOMBIA, Monte del Eden, near Haque, 9,600 feet, October; 1 ex.

Stenoma compsonaura, n. sp.

♂. 16–17 mm. Head, thorax whitish-ochreous. Palpi white, basal $\frac{3}{4}$ of second joint and base of terminal joint fuscous. Forewings rather elongate, apex rounded, termen rounded, somewhat oblique; 2 from near angle, 7 to apex (indefinite); whitish-ochreous,

all veins marked pale brownish lines; second discal stigma blackish; a marginal series of blackish dots round apex and termen: cilia whitish. Hindwings 3 and 4 connate; whitish-grey; cilia whitish.

BRAZIL, Para, July (*Parish*); FRENCH GUIANA, R. Maroni; 2 ex.

Stenoma redintegrata, n. sp.

♂. 17 mm. Head fuscous-whitish, face white. Palpi whitish, second joint suffused greyish except apex. Thorax whitish-grey-ochreous. Forewings suboblong, apex rounded-obtuse, termen rounded, slightly oblique; 2-5 approximated, 7 to termen, 8 to costa; pale fuscous, veins marked fine rather dark fuscous lines, slender whitish interneural streaks edged below with fine rather dark fuscous lines closely approximated to veins beneath them (so that there are pairs of closely adjacent dark lines); plical and second discal stigmata well-marked, black; a marginal series of black dots round apex and termen: cilia whitish. Hindwings 3 and 4 connate; pale grey, rather darker towards apex; cilia grey-whitish.

BOLIVIA, Prov. del Sara, Dept. Santa Cruz, 1500 feet; 1 ex.

Stenoma compsocharis, n. sp.

♂. 23 mm. Head, thorax white. Palpi white, a streak on basal $\frac{3}{4}$ of second joint, and extreme tip of terminal joint dark grey. Forewings somewhat dilated, apex obtuse, termen nearly straight, somewhat oblique; 2-5 approximated, 7 to termen, 8 to costa; white, somewhat sprinkled fuscous in disc; extreme edge of costa towards base and a basal dot dark fuscous; a small cloudy fuscous spot on costa at $\frac{1}{5}$, a larger one at $\frac{2}{5}$, and a still larger triangular spot at $\frac{2}{3}$ including a whitish costal dot; stigmata small, indistinct, dark fuscous, plical obliquely beyond first discal; a large fuscous dot on costa at $\frac{4}{5}$, beneath this an angulated transverse series of minute dots; a fuscous dot on costa near apex, and a terminal series of minute indistinct dots: cilia whitish, base indistinctly spotted fuscous. Hindwings 3 and 4 connate; white, apical and terminal edge interruptedly grey; cilia whitish, round apex suffused light grey.

BOLIVIA, Andes, 12,000 feet; 2 ex. Near *symmicta*, which is smaller, has grey hindwings, and dark streaks in disc of forewings.

Stenoma zephyritis, n. sp.

♂. 19 mm. Head, thorax pale ochreous, face whitish. Palpi white, basal $\frac{2}{3}$ of second joint fuscous. Forewings somewhat dilated, apex rounded-obtuse, termen rounded, little oblique; 2-5 approximated, 7 to termen, 8 to costa; pale ochreous, veins posteriorly obscurely suffused fuscous, costal edge whitish, margined fuscous beneath; stigmata blackish, plical obliquely beyond first discal; an irregular strongly curved subterminal series of dark fuscous undefined dots, deeply indented towards costa; a marginal series of dark fuscous dots round apex and termen: cilia white, basal half whitish-ochreous. Hindwings 3-5 nearly approximated

at base; ochreous-whitish; costa slenderly reflexed above from base to $\frac{2}{3}$, enclosing an ochreous-whitish hairpencil; cilia white.

BRAZIL, Santarem, August (*Parish*); 1 ex.

Stenoma crocosticta, n. sp.

♂. 23 mm. Head, thorax pale greyish-ochreous, face whitish. Palpi white, second joint with dark grey streak terminating in supramedian ring, terminal joint with base and tip blackish. Forewings hardly dilated, apex rounded, termen rounded, hardly oblique; 2-5 approximated, 7 to apex (indefinite); pale greyish-ochreous; a greyish-ferruginous dot on fold near base, and dark grey elongate dot obliquely beyond and above it; stigmata ferruginous, plical obliquely beyond first discal; a series of indistinct light ferruginous dots from beneath middle of costa very obliquely outwards to beyond second discal, then acutely angulated and continued faint and irregular to some suffusion above dorsum beyond middle; a series of cloudy subcrescentic ferruginous dots becoming dark grey towards costa, from beneath costa at $\frac{2}{5}$ very obliquely outwards strongly curved near termen and continued slightly sinuate to near dorsum about $\frac{4}{5}$; a marginal series of dark ferruginous-grey dots round posterior part of costa and termen: cilia white. Hindwings 3 and 4 connate; ochreous-whitish; cilia white, base ochreous-tinged.

PERU, Iquitos, May (*Parish*); 1 ex.

Stenoma injucunda, n. sp.

♂ ♀. 18-19 mm. Head, thorax pale greyish-ochreous, face whitish. Palpi white, second joint with dark grey streak to above middle, base and apex of terminal joint dark grey. Forewings somewhat dilated, apex obtuse, termen rounded, slightly oblique; 2 from near angle, 7 to termen, 8 to costa; light greyish-ochreous, veins suffused light fuscous; a slightly indicated fuscous dot at base of fold; stigmata blackish, plical obliquely beyond first discal; a blackish dot towards costa before middle, from beyond this a series connected by fuscous shading strongly curved round second discal stigma, indented on fold, to dorsum at $\frac{2}{3}$; a curved series of cloudy blackish dots connected by slight fuscous suffusion from costa at $\frac{4}{5}$ to dorsum before tornus, indented beneath costa, apical area beyond this suffused fuscous; a marginal series of black dots round apex and termen: cilia whitish-ochreous, tips whitish. Hindwings 3 and 4 connate; rather dark grey; cilia whitish-grey, a grey subbasal shade.

PERU, R. Napo, Jurimaguas, March, May (*Parish*); 1 ex.

Stenoma chilochema Meyr.

Hindwings ♂ with long expansible whitish hairpencil from costal edge towards base lying along costa to $\frac{2}{3}$, costal cilia thickened with scales above apex.

Stenoma antitacta, n. sp.

♂. 16 mm. Head, thorax whitish-ochreous, face whitish. Palpi whitish, basal $\frac{2}{3}$ of second joint dark grey, base and tip of

terminal joint blackish. Forewings slightly dilated, apex rounded-obtuse, termen vertical; 2 from near angle, 7 to termen, 8 to costa; whitish-ochreous slightly and irregularly sprinkled light fuscous; a small indistinct fuscous spot towards base above middle connected with dorsum by a striga; stigmata blackish, plical obliquely beyond first discal; an inwards-oblique suffused blackish spot from middle of dorsum reaching to just before plical stigma; triangular blackish spots on costa at middle and $\frac{3}{4}$, from second a strongly curved waved blackish line near termen to tornus; a marginal series of blackish dots round apical part of costa and termen: cilia whitish-ochreous. Hindwings 3 and 4 short-stalked; ochreous-whitish; cilia ochreous-whitish.

PERU, Jurimaguas, March (*Parish*); 1 ex.

Stenoma adulans, n. sp.

♂. 19 mm. Head whitish. Palpi whitish, second joint grey except tip, terminal joint suffused grey towards tip anteriorly. Thorax pale violet-grey, shoulders whitish. Forewings gradually dilated, apex obtuse, termen nearly straight, hardly oblique; 2 from near angle, 7 to termen, 8 to costa; pale violet-grey; costal edge whitish-ochreous; plical and second discal stigmata blackish-grey; an evenly curved grey line from a blackish-grey thickening on costa at $\frac{4}{5}$ near termen to tornus; a marginal series of blackish dots round apical part of costa and termen: cilia whitish. Hindwings 3 and 4 connate; rather dark grey; cilia light grey.

PERU, Jurimaguas, March (*Parish*); 1 ex.

Stenoma peridesma, n. sp.

♂. 13 mm. Head dark violet-grey, face whitish. Palpi whitish externally suffused grey. Thorax bronzy-greyish-ochreous anteriorly suffused dark grey. Forewings slightly dilated, apex rounded, termen rather obliquely rounded; 2-5 approximated, 7 to costa; light brownish-ochreous speckled grey; costal and apical edge suffused ferruginous, base of costa grey; second discal stigma small, dark grey; cilia light grey, towards tornus whitish, round apex a darker grey subbasal shade. Hindwings 3 and 4 short-stalked; grey; cilia whitish-grey, a grey subbasal shade.

BRAZIL, Obidos, September (*Parish*); 1 ex.

Stenoma volitans, n. sp.

♂. 33 mm. Head ochreous, face suffused white. Palpi whitish, second joint suffused rosy on basal $\frac{2}{5}$ and pale ochreous anteriorly, terminal joint sprinkled dark grey anteriorly. Thorax brownish-ochreous, patagia tinged purplish-grey. Forewings posteriorly dilated, costa strongly arched posteriorly, apex rounded-obtuse, termen rather obliquely rounded; 2-5 approximated, 6 to apex; ochreous-brown, on posterior third thinly strewn elongate flat pale violet-grey scales tipped blackish, a few also about fold; costal edge ochreous-rosy throughout, edged beneath greyish-violet; plical and second discal stigmata blackish: cilia light brownish-

grey. Hindwings 3 and 4 short-stalked; ochreous-yellow; cilia light yellowish tinged rosy.

PERU, Yahuas Territory; 1 ex.

Stenoma cantatrix, n. sp.

♂. 31 mm. Head, thorax whitish-fuscous speckled fuscous, face suffused whitish. Palpi whitish, second joint suffused fuscous. Forewings widest beyond middle, costa moderately arched, apex obtuse, termen somewhat oblique; 2-5 approximated, 7 to apex; whitish-fuscous closely speckled fuscous, anterior half of costa rather suffused fuscous; three nearly parallel oblique transverse dark fuscous lines, first before middle, not reaching margins, second beyond middle, becoming more oblique dorsally and ending in tornus, between first and second a parallel fascia of fuscous suffusion not reaching dorsum, third slightly excurved and meeting second on tornus; a dark fuscous dorsal line from $\frac{1}{5}$ to $\frac{3}{4}$; a suffused dark fuscous line round posterior part of costa and termen: cilia whitish-fuscous speckled fuscous. Hindwings 3 and 4 connate; rather dark grey; cilia pale grey.

BOLIVIA, Prov. del Sara, Dept. Santa Cruz, 1500 feet; 1 ex.

Stenoma argicerauna, n. sp.

♂. 23 mm. Head ochreous-whitish. Palpi whitish, second joint mixed dark fuscous and with dark fuscous subapical ring, terminal with indistinct basal and subbasal rings and anterior edge dark fuscous. Thorax pale greyish-ochreous. Forewings rather broad, broadest in middle, costa strongly arched, apex rounded off, termen rounded, rather oblique; 2-4 approximated, 6 to apparent (false) apex; pale greyish-ochreous, slightly greyer-tinged along costa towards apex; an indistinct acutely dentate whitish line running round upper margin of cell, and a transverse mark on end of cell; an almost marginal series of small indistinct whitish spots round apex and termen: cilia whitish-ochreous. Hindwings 3 and 4 very short-stalked; ochreous-whitish; cilia white.

COLOMBIA, Cantinero, 1300 feet; 1 ex. Near *algosa*.

Stenoma orneopis, n. sp.

♂. 14 mm. Head pale ochreous, crown suffused pale violet-fuscous. Palpi pale ochreous, externally suffused light violet-fuscous, terminal joint anteriorly suffused dark fuscous. Thorax violet-brownish. Forewings widest before middle, costa moderately arched, apex rounded, termen somewhat obliquely rounded; 2-5 approximated, 6 to apex (apparent); ochreous-grey-brownish, with strong violet gloss; extreme costal edge white; a rounded rosy-ochreous spot on end of cell, obscurely edged whitish; several minute whitish dots on apical margin: cilia purple-greyish. Hindwings 3 and 4 connate; blackish-grey; cilia grey, a dark grey subbasal line.

BRAZIL, Teffé, December (*Parish*); 1 ex.

Stenoma melinopa, n. sp.

♂. 18–20 mm. Head, thorax whitish-ochreous. Palpi fuscous, terminal joint white. Forewings suboblong, apex rounded off, termen rounded, vertical; 2–5 approximated, 6 to apex (apparent); light greyish-ochreous strewn dark grey dots and strigulæ; costal edge light ochreous except towards extremities; an undefined spot of grey suffusion about middle of costa; a round ochreous-greenish spot on end of cell, enclosing a more or less expressed blackish dot, beneath this is a small white dot; in one specimen an indistinct spot of greenish tinge before tornus: cilia light grey, tips paler. Hindwings 3 and 4 connate; pale greyish; cilia pale ochreous, whitish towards tips.

BRAZIL, Obidos, September (*Parish*); 2 ex.

Stenoma pratifera, n. sp.

♂. 20 mm. Head ochreous-whitish, sides of face ferruginous. Palpi whitish, second joint infuscated except apex. Thorax ochreous, a grey posterior spot. Forewings slightly broadest before middle, costa anteriorly moderately arched, apex rounded, termen rather obliquely rounded; 2 from near angle, 6 to apex (indefinite); light brown strewn grey dots and strigulæ; costal edge ochreous-yellowish; a light moss-green somewhat darker-mottled patch occupying posterior part of wing except a narrow terminal fascia of greyish suffusion, anterior edge running from costa at $\frac{2}{5}$ directly half across wing, then outwards to end of cell, and again directly to dorsum at $\frac{2}{3}$, a white dot on lower angle of cell: cilia whitish-ochreous tinged grey. Hindwings 3 and 4 connate; whitish-yellowish, basal half partially suffused pale grey; cilia whitish-yellowish, yellower round apex.

COSTA RICA; 1 ex. Near *viridis*.

Stenoma horometra, n. sp.

♂. 25 mm. Head whitish, more or less greyish on crown. Palpi whitish, second joint greyish. Thorax light grey. Forewings somewhat dilated, apex obtuse, termen slightly rounded, scarcely oblique; 2–5 closely approximated, 7 to termen, 8 to costa; whitish closely irrorated fuscous; several suffused dark brown lines, viz., one slightly curved subdorsal from base to tornus, two parallel oblique from costa before $\frac{1}{3}$ and at middle running into this, one from costa at $\frac{3}{4}$ running into second of these below middle, one rising out of this above middle, running halfway to termen, then right-angled to subdorsal line near tornus, and one round apex and termen; some fine dark brown interneural lines towards termen but not reaching it or preceding line: cilia light grey. Hindwings 3 and 4 connate; grey; cilia whitish-grey, a grey sub-basal shade.

PERU, Rio Napo, May (*Parish*); 2 ex.

Stenoma delphinodes, n. sp.

♂. 36–40 mm. Head whitish, crown more or less suffused violet-fuscous. Palpi violet-greyish, second joint with a darker

streak. Thorax violet-fuscous, a whitish posterior spot, crest posteriorly ochreous-tinged. Forewings widest about middle, costa strongly arched, apex rounded, termen rather obliquely rounded; 2-5 approximated, 6 to apex (indefinite); dark violet-fuscous, lighter and browner anteriorly; second discal stigma small, grey-whitish; a terminal series of blackish marks preceded by more or less obscure whitish suffusion: cilia fuscous, basal half on termen ochreous-whitish spotted dark fuscous, on costa dark fuscous. Hindwings 3 and 4 very short-stalked; rather dark grey, on basal half more blackish-grey; cilia grey, a darker subbasal shade.

PERU, La Oroya, Carabaya, 3000 feet, March (*Ockenden*); COLOMBIA, R. Tamaua, Choco, 320 feet, February; 3 ex.

Stenoma leucorectis, n. sp.

♂. 56-60 mm. Head, thorax, abdomen white. Palpi white, second joint light ochreous-orange anteriorly, with dark fuscous streak to above middle, terminal joint blackish towards apex. Antennal ciliations under 1. Forewings somewhat dilated, costa moderately arched, apex obtuse-pointed, termen nearly straight, oblique; 2 from near angle, 7 to apex; white: cilia white. Hindwings termen sinuate beneath apex, 3 and 4 connate; white; cilia white. Forewings and hindwings beneath with broad yellow-ochreous costal bands, sometimes some grey suffusion at apex of forewings.

BRAZIL, Para, July (*Parish*); and Leopoldina, Espiritu Santo; 2 ex.

Stenoma auxoleuca, n. sp.

♂ ♀. 47-53 mm. Head, thorax, abdomen white. Palpi white, second joint orange anteriorly, a blackish streak on basal half, tip of terminal joint blackish-grey. Antennal ciliations ♂ $1\frac{1}{2}$. Forewings somewhat dilated, costa gently arched, apex rounded-obtuse, termen rounded, rather oblique; 2 from towards angle, 7 to apex (indefinite); white: cilia white. Hindwings 3 and 4 connate; white; cilia white. Forewings beneath suffused grey-yellowish towards costa and tinged grey-yellowish on cell and costal half, but leaving a slender white median longitudinal streak, and a white fascia round apex and termen; hindwings tinged grey-yellowish towards costa.

BRAZIL, Leopoldina, Espiritu Santo; 2 ex.

Stenoma maturescens, n. sp.

♂ ♀. 42-50 mm. Head, thorax, abdomen white. Palpi white, second joint more or less orange, a dark fuscous streak on basal $\frac{1}{3}$ - $\frac{2}{3}$, terminal joint usually blackish towards tips or sometimes wholly blackish or wholly white. Antennal ciliations ♂ $1\frac{1}{2}$. Forewings dilated, costa moderately arched, apex rounded, termen rounded, rather oblique; 2 from towards angle, 7 to apex (indefinite); white: cilia white. Hindwings termen rounded, 3 and 4 connate; white; cilia white. Forewings beneath suffused light ochreous-yellowish except towards dorsum, deeper yellow-ochreous towards

costa and apex, round apical edge more or less suffused fuscous; hindwings beneath suffused yellow-ochreous towards costa.

FRENCH GUIANA, R. Maroni; 10 ex.

Stenoma argonais, n. sp.

♂. 42 mm. Head, thorax, abdomen white. Palpi white, second joint pale orange anteriorly, a dark fuscous streak on basal half, terminal joint blackish-grey towards apex. Antennal ciliations under 1. Forewings dilated, costa gently arched, apex rounded-obtuse, termen rounded, rather oblique; 2 from towards angle, 7 to apex (indefinite); white: cilia white. Hindwings 3 and 4 connate; white; cilia white. Forewings beneath tinged yellowish except towards dorsum, towards costa and apex suffused deeper yellow-ochreous, apical edge suffused fuscous; hindwings beneath suffused ochreous-yellowish towards costa and apical part of termen.

BRAZIL, Fonte Boa, February (*Parish*); 1 ex.

Stenoma megaleuca Meyr.

Thorax and abdomen of original type (still unique) described as white (I supposed them to be discoloured); actually the thorax is tinged greyish-ochreous except patagia, abdomen suffused pale greyish-ochreous becoming greyer posteriorly; I now think this colouring may be natural, but am not certain; there is nothing of the sort in any of the allied species. Otherwise the species is nearest *auxoleuca*.

Stenoma isarga, n. sp.

♀. 40 mm. Head, thorax, abdomen white. Palpi white, basal half of second joint orange-ochreous with a blackish streak, terminal joint blackish except towards base. Middle and posterior legs wholly white. Forewings rather dilated, costa gently arched, apex rounded-obtuse, termen rounded, rather oblique; white: cilia white. Hindwings 3 and 4 connate; white; cilia white. Wings beneath wholly white.

BOLIVIA, Prov. del Sara, Dept. Santa Cruz, 1500 feet, November; 1 ex.

Stenoma philomela, n. sp.

♂. 22 mm. Head, thorax, abdomen white. Palpi white, second joint ochreous-orange anteriorly, a blackish streak on basal $\frac{2}{3}$, tip of terminal joint blackish-grey. Antennal ciliations 1. Forewings dilated, costa moderately arched, apex rounded, termen rounded, rather oblique; 2 from near angle, 7 to apex (indefinite); white: cilia white. Hindwings 3 and 4 connate; yellow-whitish-grey, apical edge yellow; cilia white. Forewings beneath suffused light yellowish except towards dorsum, towards costa suffused ochreous-orange, apical edge suffused dark grey; hindwings beneath suffused yellowish towards costa.

PERU, Jurimaguas, March (*Parish*); 1 ex.

Stenoma subita, n. sp.

♂. 26 mm. Head, thorax rather dark grey. Palpi grey. Forewings elongate, apex rounded, termen rounded, somewhat